

## **Table of Contents**

Overview	3
Summary	3
Describing the Survey Process	
Sampling Plan	
Mailing of Questionnaires	3
Inclusion/Exclusion Criteria	
Developing the Indicators	5
The Four Indicators reported in Hospital Report 2007: Emergency Department Care	e5
The Seven Picker-based Indicators	6
The Two Functional Indicators	8
Calculating the Indicator Scores	10
Risk-Adjustment	11
Risk-Adjustment Procedure	12
Risk-adjustment for Sex-Disaggregated Patient Satisfaction Results	13
Missing Values for Predictors Used in Risk-Adjustment	13
Descriptive Statistics for Predictors Used in Risk-Adjustment	13
Patient-response Statistics	13
Proxy-response Statistics	14
Weighting	15
Hospital Site Weights	15
Corporation Weights	
Response Rates	16
Hospital-Specific Analysis: Assessing Relative Performance	16
Confidence Intervals and the Allocation Bias	
Statistics Provided for Hospital Comparisons with Provincial Results	20
Appendix A: Questionnaire Items	24

## **Overview**

Patient Satisfaction monitors the patient's perceptions of the hospital care they received. Hospital Report '99 was the first report that allowed for province-wide comparisons among Ontario hospitals in terms of levels of patient satisfaction. Hospital Report 2001, 2002, 2003, 2005, 2006 then built on these province-wide comparisons and presented levels of patient satisfaction for each participating Ontario hospital. Hospital Report 2007: Emergency Department builds on the substantial research documenting what is important to patients, how patients evaluate care, and how to measure this. Even though there are generally no standards for levels of satisfaction that are desired or necessary in Canadian hospitals, knowledge gained from patient satisfaction surveys can set a direction for quality improvement in hospitals for outcomes that are important to patients.

This year's patient satisfaction results reflect data from April 1, 2005 to March 31, 2006. There will be no half data years reported in this year's report. Results are provided online in the e-Scorecard and in the Executive Summary.

## **Summary**

Approximately 115,000 individuals from 90 participating hospitals in Ontario were sampled for the 2005-2006 Emergency Department Care Patient Satisfaction analysis. Patients discharged between April 2005 and March 2006 who met the inclusion criteria were included in the sample. Approximately 31% of the sampled individuals returned their questionnaires.

## **Describing the Survey Process**

## Sampling Plan

Each participating hospital corporation and NRC+Picker collaboratively established a sampling plan. Deciding factors influencing the agreed-upon sampling plan included budget, achieving reasonable response rates, and which sites within the corporation were of primary interest. A minimum of 100 valid survey responses is required for a hospital's results to be displayed in the publicly released summary report.

Hospitals were then charged with the responsibility of sending patient data files to NRC+Picker every month. For *Hospital Report 2007: Emergency Department Care*, data was collected for all 12 months of the 2005-2006 fiscal year. Then, according to each hospital's sampling plan, a random sample was drawn from the patient data files, and surveys were mailed.

Questionnaires were not sent to deceased patients, psychiatric patients, infants less than 10 days old, patients with no fixed address, or patients who presented with sexual assault or other sensitive issues.

## Mailing of Questionnaires

Included in each patient mailing was an explanatory cover letter, a return envelope (postage-paid), and the questionnaire itself. The first mailing went out within a couple of weeks of NRC's reception of a hospital's monthly patient data file. To increase response rates, there was a second wave of mailings to patients whose first questionnaires were not returned within three weeks of the original mailing date.

Approximately, 114,666 surveys were sent in 2005-2006, yielding 113,071 returns. Of these, 34,984 (approx. 31% of all valid returns, and 30.5% of all surveys) met all inclusion criteria, and are included in the analysis.

#### Inclusion/Exclusion Criteria

Surveys that were returned, but that did not have a single valid response, were treated as non-responses and dropped from the analysis.

Surveys that were returned, but that did not have a single valid response, were treated as non-responses and dropped from the analysis. If a record had no valid responses to any of the evaluative questions on the questionnaire (i.e. only had responses to demographic-type questions), then it was seen as having insufficient data and was excluded from the subsequent analysis.

Finally, for a record to be included in the analysis, it had to have valid responses for at least half of the evaluative questions that were used to build the indicators. This amounts to 17 valid responses out of the 34 evaluative questions that make up the four indicators that are included in the Executive Summary report. See Appendix A for a list of the questionnaire items. Note that it is at this stage that the volumes were calculated to determine whether a hospital passed the 100-case volume screen.

## **Developing the Indicators**

Thirty-four of the 58 Emergency Department Care survey questions were combined to form four Hospital Report 2007 ED indicators as well as seven Picker-based indicators and two functional indicators. All ED patient satisfaction indicators are available to participating hospitals in the e-Scorecard (www.hospitalreport.ca).

Expert opinion, from patient focus groups and hospital staff surveys, and discussions among team members lead to the amalgamation of NRC groupings and items into four indicators. This took place in two steps: NRC groupings were first adjusted by moving some items and combining some categories; then this reduced set of indicators was further reduced into four categories.

## The Four Indicators reported in Hospital Report 2007: Emergency Department Care

Traditionally, these four indicators were reported in previous Hospital Report executive summaries and e-Scorecard.

**1. Overall Impressions:** A patient's assessment, overall, of their hospital stay. Based on 4 survey questions.

#### **Survey Questions:**

- Did you have confidence and trust in the doctors treating you?
- Did you have confidence and trust in the nurses treating you?
- Overall, how would you rate the care you received at the hospital?
- Would you recommend this hospital to your friends and family?
- Communication: A patient's assessment of how well information was communicated to them or their family during their ED stay. Based on 14 survey questions.

#### **Survey Questions:**

- When you arrived at the Emergency Department, did the first person who took your information answer your questions?
- If you had to wait to be seen, did someone from the Emergency Department explain the reason for the delay?
- Did someone in the Emergency Department help get your messages to family or friends?
- When you had important questions to ask a doctor, did you get answers you could understand?
- If you had any anxieties or fears about your condition or treatment, did a doctor discuss them with you?
- When you had important questions to ask a nurse, did you get answers you could understand?
- If you had any anxieties or fears about your condition or treatment, did a nurse discuss them with you?
- Did someone explain why you needed these tests in a way that you could understand?

- Did someone explain the results of the tests in a way that you could understand?
- Did they tell you what danger signals about your illness or operation to watch for after you went home?
- Did someone explain how to take the new medications?
- Did someone tell you about medication side effects to watch for when you went home?
- Did you know who to call if you needed help or had more questions after you left the hospital?
- How would you rate the explanation of what was done to you?
- **3. Consideration:** A patient's assessment of whether they were treated with consideration. Based on 6 survey questions.

#### **Survey Questions:**

- How would you rate the courtesy of the first person who took your information?
- How would you rate the courtesy of your doctors?
- How would you rate the courtesy of your nurses?
- Did each hospital staff person treat you with dignity and respect?
- Did you have enough say about your care?
- How would you rate the courtesy of the Emergency Department staff?
- **4. Responsiveness**: A patient's assessment of the organization and responsiveness to their needs during their ED stay. Based on 10 survey questions.

#### **Survey Questions:**

- Did you have to wait too long to see a doctor?
- Did you wait too long for this other doctor or specialist?
- How would you rate the availability of your nurses?
- Did you wait too long to get your tests?
- Do you think that the Emergency Department staff did everything they could to help control pain?
- Overall, how much pain medicine did you get?
- How would you rate the amount of time you spent in the Emergency Department?
- While you were in the Emergency Department, were you able to get all the services you needed?
- While you were in the Emergency Department, were there times when you did not get the help you needed?
- How would you rate how well the doctors and nurses worked together?

#### The Seven Picker-based Indicators

These Picker-based indicators are only available in the e-Scorecard.

#### 1. Overall Satisfaction

#### **Survey Questions:**

- How would you rate the amount of time you spent in the Emergency Department?
- How would you rate the explanation of what was done to you?
- How would you rate how well the doctors and nurses worked together?
- Was the entire Emergency Department as clean as it should have been?
- Would you recommend this Emergency Department to family and friends?

#### 2. Coordination of Care and Access

#### **Survey Questions:**

- While you were in the Emergency Department, were there times when you did not get the help you needed?
- While you were in the Emergency Department, were you able to get all the services you needed?
- Did you have to wait too long to see a doctor?
- Did you wait too long for this other doctor or specialist?
- How would you rate the availability of your nurses?
- Did you wait too long to get your tests?
- After you arrived at the Emergency Department, how long was it until you talked to a NURSE about your illness or injury?
- Was there one particular doctor in charge of your care in the Emergency Department?

#### 3. Physical Comfort

#### **Survey Questions:**

- Do you think that the Emergency Department staff did everything they could to help control your pain?
- · Overall, how much pain medicine did you get?

#### 4. Respect for Patient Preferences and Courtesy

#### **Survey Questions:**

- Did each hospital staff person treat you with dignity and respect?
- Did you have enough say about your care?
- Did you feel you had enough privacy during your Emergency Department visit?
- How would you rate the courtesy of your doctors?
- How would you rate the courtesy of your nurses?
- How would you rate the courtesy of the first person who took your information?
- How would you rate the courtesy of the Emergency Department staff?

#### 5. Information and Education

#### **Survey Questions:**

 When you had important questions to ask a nurse, did you get answers you could understand?

- When you had important questions to ask a doctor, did you get answers you could understand?
- Did someone explain the results of the tests in a way that you could understand?
- Were the possible causes of your problem explained in a way that you could understand?
- Did someone explain why you needed these tests in a way that you could understand?
- When you arrived at the Emergency Department, did the first person who took your information answer your questions?
- If you had to wait to be seen, did someone from the Emergency Department explain the reason for the delay?

#### 6. Continuity and Transition

#### **Survey Questions:**

- Did someone explain how to take the new medications?
- Did you know who to call if you needed help or had more questions after you left the Emergency Department?
- Did someone tell you about side effects the medicines might have?
- Were you told what danger signals about your illness or injury to watch out for when you got home?

#### 7. Emotional Support

#### **Survey Questions:**

- If you had any anxieties or fears about your condition or treatment, did a doctor discuss them with you?
- Did you have confidence and trust in the nurses treating you?
- Did you have confidence and trust in the doctors treating you?
- If you had any anxieties or fears about your condition or treatment, did a nurse discuss them with you?
- Did someone in the Emergency Department help get your messages to family or friends?

#### The Two Functional Indicators

These indicators are assessments of the quality of the care received from health care providers. These indicators are only available via the e-Scorecard.

#### 1. Physician Care

#### **Survey Questions:**

- When you had important questions to ask a doctor, did you get answers you could understand?
- If you had any anxieties or fears about your condition or treatment, did a doctor discuss them with you?
- Did you have confidence and trust in the doctors treating you?

- Did you wait too long for this other doctor or specialist?
- How would you rate the courtesy of your doctors?

#### 2. Nursing Care

#### **Survey Questions:**

- When you had important questions to ask a nurse, did you get answers you could understand?
- If you had any anxieties or fears about your condition or treatment, did a nurse discuss them with you?
- Did you have confidence and trust in the nurses treating you?
- How would you rate the courtesy of your nurses?
- How would you rate the availability of your nurses?

#### Calculating the Indicator Scores

Each question within an indicator carries equal weighting in the calculation of the indicator score. Indicator scores are calculated at the patient-level, where a patient has valid, score-able responses for at least half of the items belonging to that indicator.

A variety of response scales were used in the patient satisfaction questionnaire. Some questions employed a Likert-type scale with five response choices: "Poor", "Fair", "Good", "Very Good", "Excellent". These were assigned the following scores: Poor = 0, Fair = 25, Good = 50, Very Good = 75, Excellent = 100.

Other questions used a three point scale with responses resembling: "Yes, Always", "Yes, Sometimes", "No". These were assigned the following scores: "Yes, Always" = 100, "Yes, Sometimes" = 50, "No" = 0. A number of these questions had a fourth viable selection. For example, the question "Did someone tell you about side effects the medicines might have?" had the response options "Didn't need explanation". This was an acceptable response to the question, but was not assigned a score.

There were also questions using three-point scoring schemes with responses similar to those outlined above, but where the awarding of points was reversed. For example, the question "Did you have to wait too long to see a doctor?" would be scored: "Yes, definitely" = 0, "Yes, somewhat" = 50, "No" = 100.

For a full listing of the questionnaire's items and related scoring, see Appendix A.

## **Risk-Adjustment**

In comparing hospitals on the quality of their patient care, it is important to take into account differences in patient characteristics that may vary systematically among hospitals. In clinical research this is called risk-adjustment, where patient scores are adjusted to remove pre-existing influences. This issue is particularly important because certain groups of patients tend to systematically report lower levels of satisfaction than other groups. If a hospital tends to serve a disproportionate number of such patients, it may be unfairly reported as having lower patient satisfaction, when in fact, satisfaction may be comparable to another hospital with higher satisfaction scores that simply serves a different population. Therefore, to improve hospital comparability, a statistical technique called hierarchical modeling was used to control for possible differences in pre-existing patient characteristics. Hierarchical modeling is a way of looking at outcomes, in this case patient satisfaction scores, and their relationships to particular factors that can be used to predict them. Hierarchical models differ from multivariate regression models (used in past years of the Hospital Report) in that they take into account the fact that the data are nested into distinct groups (in this case, patients nested into hospital corporations). Hierarchical models allow the hospital corporation's effect on the resulting model to vary from hospital to hospital, essentially treating the corporation itself as a risk-adjustment variable. As in previous Reports, five different factors, or predictor variables, were used at some stage of the risk-adjustment.

Characteristics that are most commonly used in risk-adjustment of patient satisfaction scores are age and gender<sup>6,7</sup>. Previous research has found that older patients tend to report slightly higher satisfaction than younger patients. Similarly, men tend to be slightly more satisfied with their care than women. Both of these variables were used in the risk-adjustment.

In addition, questions assessing patients' perception of the severity of their conditions were also used. Less healthy patients may systematically report different levels (higher or lower) of satisfaction than do relatively healthier patients, regardless of the quality of hospital care they received<sup>8</sup>. Thus, patients' self-assessed health status was used in the risk-adjustment. Results were also adjusted for whether the patient had a regular family physician/general practitioner.

Finally, a question asking if someone other than the patient had completed the survey was considered. A hospital might care for relatively more patients who were less able to complete the survey because of age, language barriers, or illness, for example. If these proxy respondents tended to give different (higher or lower) responses than actual patients, then a biased representation of the satisfaction scores for that hospital would result. A decision was made not to include the proxy variable in the risk-adjustment model, but rather, to divide the dataset into 'patient' responses and 'proxy' responses, and then to risk-adjust them separately.

The risk-adjustment was performed separately for each of the 13 indicators (four Hospital Report indicators, seven Picker-based indicators, two Functional indicators). The results of this analysis were then used to construct new patient satisfaction scores, removing any variation in the original scores that was attributable to the predictors used in the risk-adjustment.

## Risk-Adjustment Procedure

- The patient satisfaction data was divided into two sets: one including surveys that were completed by the patients themselves; the other including surveys that were completed by someone else (i.e. by proxy).
- The model that was chosen to risk-adjust the 'patient' responses included the fixed effects of the variables age, gender, self-assessed health, whether the patient had a regular family physician, and a squared age term (age<sup>2</sup>). This model also included the random effects of the hospital corporation (allowing the intercept of the model to vary by corporation) and age (allowing the effect on the model of an individual's age to vary by corporation). The model that was chosen to risk-adjust the 'proxy' responses included the variables self-assessed health, and whether the patient had a regular family physician. Again, this model included the random effect of the hospital corporation.
- Based on the resulting models, residual scores were calculated for each patient level record. The residual is the portion of the original score that cannot be explained or predicted by the model. The residual score is, therefore, the result of some unknown source of variation in the data. Note: a patient's 'residual' score plus their 'predicted' score adds to their original score. An individual's 'predicted' score is the score that is output when all of the patient's risk-adjustment characteristics are put into the model.
- Each patient's residual score is added to the overall mean of the 'predicted' scores to yield their risk-adjusted score.
- Before re-combining the newly risk-adjusted 'patient' and 'proxy' datasets, a
   'Proxy bump' is applied. The mean patient scores for each of these datasets
   (patient, proxy) were calculated by approximate 10-year age bands (18-29, 3039, etc.). The difference between the mean scores of the corresponding age
   bands in the two datasets was calculated and added to the 'proxy' responses to
   bump them to the same level as the 'patient' responses (the 'proxy' means were
   lower than the 'patient' means in all but one instance, where no bump was
   applied).
- The 'patient' and 'proxy' datasets are then combined.

Note: Missing values in the risk-adjustment variables were assigned alternative valid responses (see Tables 1a & 1b).

# Risk-adjustment for Sex-Disaggregated Patient Satisfaction Results

The Patient Satisfaction Sex-Disaggregated results were risk-adjusted slightly differently from those in the regular analysis. So as not to compensate for the differences between males and females, the risk-adjustment was performed on the data using the same methodology as outlined above, but without controlling for the effect of gender. As a result, variation in the data due to differences between males and females will remain intact.

All other aspects of the risk-adjustment for the sex-disaggregated results – including the separate adjustments for 'patient' and 'proxy' groups, the other covariates in the models, and the 'proxy bump' – are consistent with the methodologies of the regular patient satisfaction analysis.

## Missing Values for Predictors Used in Risk-Adjustment

Table 1: FY 2005/2006

Predictor	Frequency	Percent	Default substitution for missing
			values
Proxy Question	761	2.23%	A proxy response
Age	462	1.35%	N/A
Gender	0	0%	Female
Self-assessed health	542	1.59%	The mode for all other patients from
			the same corporation
Whether patient has	387	1.13%	No regular family physician
regular family			
physician			

## Descriptive Statistics for Predictors Used in Risk-Adjustment

Table 2: Descriptive Statistics for Proxy Question FY 2005/2006

	Frequency	Percent
Patient response	25470	74.69%
Proxy response	8629	25.31%

## Patient-response Statistics

Table 3: Descriptive Statistics for Age FY 2005/2006

N	Min.	Max.	Mean	Std Dev.	Skewness	Kurtosis
25470	16	101	52.38	18.64	-0.10	-0.89

Table 4: Descriptive Statistics for Gender FY 2005/2006

Table 4. Descriptive diatistics for deflact 1.1. 2000/2000				
	Frequency	Percent		
Female	14621	57.40		
Male	10849	42.60		

Table 5: Descriptive Statistics for Self-Assessed Health FY 2005/2006

	Frequency	Percent
Poor	1027	4.03%
Fair	4071	15.98%
Good	8983	35.27%
Very Good	8100	31.80%
Excellent	3289	12.91%

Table 6: Descriptive Statistics for whether patient has a regular family physician FY 2005/2006

	Frequency	Percent
Yes	23241	91.25%
No	2229	8.75%

## **Proxy-response Statistics**

Table 7: Descriptive Statistics for Age FY 2005/2006

N	Min.	Max.	Mean	Std Dev.	<b>Skewness</b>	Kurtosis
8629	0	101	26.02	29.46	1.01	-0.59

Table 8: Descriptive Statistics for Gender FY 2005/2006

	Frequency	Percent
Female	3960	45.89%
Male	4669	54.11%

Table 9: Descriptive Statistics for Self-Assessed Health FY 2005/2006

	Frequency	Percent
Poor	465	5.39%
Fair	923	10.70%
Good	2143	24.83%
Very Good	2895	33.55%
Excellent	2203	25.53%

Table 10: Descriptive Statistics whether patient has a regular family physician FY 2005/2006

	Frequency	Percent	
Yes	7966	92.32%	
No	663	7.68%	

## Weighting

## Hospital Site Weights

Weights are necessary in hospital-specific analysis to ensure that a hospital's patient sample reflects the actual discharge pattern of the units/sites that make up that hospital corporation.

Hospital site weights were calculated using the patient data files that hospitals provided to NRC. The expectation was that these patient data files would yield a reasonable approximation of actual discharge patterns from a hospital. Within each hospital corporation, a weight was calculated for each site and month by comparing the sample population to the discharge population. This means that each site within a corporation can have as many as 12 distinct weights (one for each month of data). For every corporation/site/month combination where it was not possible to calculate a weight, a weight of 1 was assigned.

For each hospital corporation, the goal of weighting is to make the distribution of the sampled population resemble that of the discharged population. Ideally, site X would represent the same proportion of cases in the sample population as it does in the discharge population. If this is the case, then a weight of 1 is assigned to each case within site X. If site X is under-represented in the sample (i.e. it has proportionately fewer cases in the sample population than in the discharge population), than every case from that site will receive a weight greater than one to compensate. Similarly, cases from sites that are over-represented will receive weights smaller than one (but greater than zero; there are no negative weights).

These hospital site weights were applied throughout the patient satisfaction analysis. They were applied in the risk adjustment with hierarchical models, and the calculation of hospital-specific indicator scores. These were also used, along with corporation weights, in calculating indicator scores by region (LHIN) and peer group, and in calculating province-level satisfaction scores (note: these provincial scores are not those used for the performance allocation).

## **Corporation Weights**

A different set of weights was required (along with the hospital unit weights) for the calculation of indicator scores at the peer group, regional, and provincial level. Where above, we discussed weights for sites within a corporation, here we need to consider weights for corporations within the province. In calculating broader indicator scores, we have to ensure that the size of a corporation's sample population within the entire sample is proportional to the size of its discharge population within the entire discharge population (i.e. all discharges for the participating Ontario hospitals). A corporation that is either over-represented or under-represented in the sample will see weights applied to its cases accordingly. Discharge information from NRC+Picker's sampling plan was used to calculate the corporation weights.

## **Response Rates**

For the 2005-2006 fiscal year, the overall response rate was 30%. Females had a response rate of 31.9% while males had a response rate of 28.1%. The mean hospital response rate was 31.2%, and the median response rate was 31.1%. The lowest response rate for a given hospital corporation was 20.2%.

# Hospital-Specific Analysis: Assessing Relative Performance


Each hospital's performance on each indicator is designated "above average", "average", or "below average". For the patient satisfaction indicators, two criteria were used to assess each hospital's performance. Statistical significance was used to assign hospitals to the above average performance category. This criterion alone, however, was insufficient when designating hospitals as having below average performance. The calculation of statistical significance relies heavily on sample size, and different hospitals in this project had dramatically different sample sizes. It was possible to find statistically significant differences from average in hospitals with a larger sample size, even though those differences were small and not substantive. A second criterion, described below, was used for assigning hospitals to the below average performance category.

A 99.9% Confidence Interval (CI) was calculated for each indicator score within each hospital. Each hospital was compared to the average indicator score for all hospitals (the grand mean):

- "Above average" performance allocations were given to hospitals with indicator scores that were significantly statistically above the grand mean, using the 99.9% CI. This means that the lower bound of the confidence interval was higher than the grand mean.
- "Provincial Average" performance allocations were given to hospitals with indicator scores not significantly different from the grand mean.
- "Below average" performance allocations were given to hospitals with indicator scores both statistically significantly below the grand mean and lower than all nonsignificant hospital indicator scores. This means that the upper bound of the confidence interval was below the grand mean, and the hospital's indicator score was lower than the scores of all hospitals that were allocated an "average" performance.

#### Step 1: Calculate 99.9% Cls and Compare to Grand Mean

Figure 1.1: Procedure for Determining a Hospital's Performance Category



**HOSPITAL A:** The grand mean is above the upper 99.9% CI. The average indicator score for hospital A is estimated to be less than the grand mean (average of all hospital average) 999 times out of 1000.

ALLOCATION: Below average

**HOSPITAL B:** The grand mean falls between the upper and lower 99.9% CI of the hospital average. The average indicator score for hospital B is estimated to not be significantly different than the grand mean 999 times out of 1000.


ALLOCATION: Provincial Average


**HOSPITAL C:** The grand mean is below the lower 99.9% CI. The average indicator score for hospital C is estimated to be greater than the grand mean 999 times out of 1000.

ALLOCATION: Above average

#### Step 2: Apply Low Cut-Off Point

Figure 1.2: Applying the Low Cut-Off Point for Performance Allocations





#### Confidence Intervals and the Allocation Bias

The value for each patient satisfaction indicator represents the average response of patients surveyed. Hospitals that survey a greater number of patients tend to have a narrower confidence interval. This means that we are generally more confident (99.9%) that the true indicator score is within this narrow band. Conversely, hospitals that survey fewer patients would be more likely to have a wider confidence interval. Therefore, hospitals with an average score below the grand mean would be more likely to receive an allocation of "provincial average" when they have a wider confidence interval while a hospital with the same score with a narrow confidence interval would be more likely to receive an allocation of "below average". To prevent this problem, a low cut-off point was used for the "below average" allocation.

#### 99.9% Low Cut-off Point (HOSPITAL D & E)

Hospital D has the same average score as hospital E. However, based on the allocation process previously described, hospital D would receive a lower allocation (below average) than hospital E (provincial average) due to the size of the confidence intervals. A low cut-off point is applied to prevent this bias against hospitals that have a greater sample size/narrower confidence interval.

The 99.9% low cut-off point is determined by identifying the lowest indicator score of those hospitals that received an allocation of "provincial average" (and for which the grand mean fell within the 99.9% CI). Any hospital with an average score which falls between this low cut-off point and the grand mean receives an allocation of "provincial average" irrespective of whether the upper 99.9% confidence limit falls below the "provincial average". Therefore, hospital D which has the same average score as hospital E would also receive an allocation of "provincial average".

#### What About an Above Average Cut-off Point? (HOSPITAL F & G)

Note that although a low cut-off point was applied to prevent a bias against poorer allocation of performance for hospitals with a greater sample size and narrower confidence interval, this process was not applied for above average performance. Conversely applied, this process would penalize hospitals that have a greater sample size and narrower confidence interval by lowering the performance score when in fact the hospital is significantly above average. As a result, while both hospital F and G have the same average score, hospital G would receive an "above average" allocation, while hospital F would receive a "provincial average" allocation.

The following table represents the lower cut off points for all the indicators.

Table 11: 99.9% Low Cut-Offs (FY 2005-2006 only)

Indicator Group	Indicator	99.9% Low Cut-Off
Hospital Report Indicators	Overall Impressions	71.4864
	Responsiveness	62.4827
	Consideration	70.9886
	Communication	62.3951
Picker-based Indicators	Overall Satisfaction	64.6352
	Coordination of Care and	64.2916
	Access	
	Physical Comfort	57.6521
	Respect for Patient	71.4150
	Preferences and Courtesy	
	Information and Education	68.0425
	Continuity and Transition	55.3276
	Emotional Support	67.6736
Functional Indicators	Physician Care	70.5630
	Nursing Care	68.2675

# Statistics Provided for Hospital Comparisons with Provincial Results

Descriptive information is provided on the following pages. This will allow individual hospital corporations to be compared with the province as a whole. Included for each scale for all hospitals overall, and for hospitals by peer group, is the valid N, the grand mean, the standard deviation of the hospital averages, the median, and the range. Note: The "All Hospitals" and peer group means presented in these tables are calculated using weighted and risk-adjusted data, and they exclude data from participating hospitals that did not pass the volume screen. For these reasons, the means will not match those presented in the Executive Summary or E-scorecard. They are presented as a useful tool for hospital comparison.

#### **HOSPITAL REPORT INDICATORS:**

Table 12: Overall Impressions FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	74.67	82.87	73.54	75.58
Std Deviation	24.57	14.93	25.17	27.64
Minimum	61.19	76.60	61.19	68.51
25 <sup>th</sup> Percentile	73.64	81.67	71.49	74.05
Median	76.51	83.49	75.24	75.91
75 <sup>th</sup> Percentile	80.80	86.55	78.15	78.51
Maximum	89.38	89.38	85.90	82.68

Table 13: Responsiveness FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	65.70	77.28	64.67	65.00
Std Deviation	24.09	14.60	24.51	27.17
Minimum	51.63	68.48	51.63	59.36
25 <sup>th</sup> Percentile	64.36	75.11	62.92	61.62
Median	67.38	77.05	66.64	65.17
75 <sup>th</sup> Percentile	73.65	81.19	69.13	65.88
Maximum	85.61	85.61	80.45	73.19

Table 14: Consideration FY 2005/2006

14510 111 9011014014110111 1 2000/2000					
	All Hospitals	Small	Community	Teaching	
Valid N	90	18	58	14	
Mean	72.98	80.09	72.11	73.37	
Std Deviation	21.18	13.46	21.56	24.13	
Minimum	60.86	72.63	60.86	67.82	
25 <sup>th</sup> Percentile	71.85	79.15	70.99	71.39	

Median	74.32	80.87	73.59	73.83
75 <sup>th</sup> Percentile	78.17	83.22	76.12	75.75
Maximum	85.41	85.41	83.43	80.86

Table 15: Communication FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	67.25	74.16	66.37	67.95
Std Deviation	24.17	16.89	24.60	26.56
Minimum	57.35	62.40	57.35	62.65
25 <sup>th</sup> Percentile	66.08	72.02	65.76	64.87
Median	68.13	76.51	67.42	67.85
75 <sup>th</sup> Percentile	71.63	79.46	69.69	70.30
Maximum	80.35	80.35	78.63	72.48

#### PICKER-BASED INDICATORS:

Table 16: Overall Satisfaction FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	66.13	76.48	64.90	66.61
Std Deviation	23.26	14.45	23.73	25.91
Minimum	52.46	68.04	52.46	61.44
25 <sup>th</sup> Percentile	64.64	74.69	62.79	64.64
Median	68.06	77.29	66.63	66.27
75 <sup>th</sup> Percentile	73.81	80.80	71.03	69.73
Maximum	84.80	84.80	79.66	73.81

Table 17: Coordination of Care and Access FY 2005/2006

Table 17: Goordination of Gare and Access 11 2000/2000					
	All Hospitals	Small	Community	Teaching	
Valid N	90	18	58	14	
Mean	65.50	77.16	64.80	63.55	
Std Deviation	23.34	14.02	23.74	26.28	
Minimum	50.75	68.88	50.75	57.01	
25 <sup>th</sup> Percentile	63.20	75.15	62.73	59.78	
Median	67.73	77.39	67.51	63.82	
75 <sup>th</sup> Percentile	73.28	80.75	69.22	65.97	
Maximum	85.00	85.00	80.75	70.37	

Table 18: Physical Comfort FY 2005/2006

	<b>All Hospitals</b>	Small	Community	Teaching
Valid N	90	18	58	14
Mean	65.90	74.89	64.89	66.59
Std Deviation	38.72	26.55	39.05	44.09
Minimum	52.25	63.83	52.25	61.26

25 <sup>th</sup> Percentile	63.58	74.03	62.41	62.77
Median	68.00	76.37	67.52	65.16
75 <sup>th</sup> Percentile	72.52	78.74	69.39	68.64
Maximum	82.56	82.56	77.47	74.91

Table 19: Respect for Patient Preferences and Courtesy FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	73.85	81.10	73.01	74.09
Std Deviation	20.78	13.04	21.18	23.61
Minimum	61.04	73.81	61.04	69.04
25 <sup>th</sup> Percentile	72.63	79.98	72.12	71.80
Median	75.56	82.06	74.64	74.71
75 <sup>th</sup> Percentile	79.37	83.97	77.02	76.36
Maximum	86.82	86.82	84.43	81.28

Table 20: Information and Education FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	72.09	79.02	71.25	72.73
Std Deviation	24.15	16.16	24.61	26.60
Minimum	63.55	68.49	63.55	67.87
25 <sup>th</sup> Percentile	70.79	75.98	70.17	70.98
Median	73.28	80.12	72.14	72.56
75 <sup>th</sup> Percentile	75.93	83.78	74.55	75.29
Maximum	85.34	85.34	82.10	75.93

Table 21: Continuity and Transition FY 2005/2006

	All Hospitals	Small	Community	Teaching
Valid N	90	18	58	14
Mean	61.15	68.77	60.20	61.65
Std Deviation	34.19	24.30	34.69	38.59
Minimum	51.48	56.64	51.48	57.29
25 <sup>th</sup> Percentile	59.59	65.77	58.31	59.51
Median	62.18	71.87	61.45	60.64
75 <sup>th</sup> Percentile	65.88	74.01	64.58	64.40
Maximum	74.90	74.63	74.90	69.98

Table 23: Emotional Support FY 2005/2006

Table 201 Elliotichal Cappett 1 1 2000/2000					
	All Hospitals	Small	Community	Teaching	
Valid N	90	18	58	14	
Mean	72.90	80.33	72.08	73.31	
Std Deviation	27.00	17.60	27.51	29.83	
Minimum	58.89	72.78	58.89	66.89	
25 <sup>th</sup> Percentile	72.31	79.14	71.43	70.35	

Median	74.57	81.18	73.71	73.59
75 <sup>th</sup> Percentile	78.52	84.37	76.38	75.47
Maximum	85.72	85.72	83.82	80.13

#### **FUNCTIONAL INDICATORS:**

Table 24: Physician Care FY 2005/2006

144010 2 11 1 11 y 0101411 0 410 1 1 2000/2000					
	All Hospitals	Small	Community	Teaching	
Valid N	90	18	58	14	
Mean	65.90	74.89	64.89	66.59	
Std Deviation	38.72	26.55	39.05	44.09	
Minimum	52.25	63.83	52.25	61.26	
25 <sup>th</sup> Percentile	63.58	74.03	62.41	62.77	
Median	68.00	76.37	67.52	65.16	
75 <sup>th</sup> Percentile	72.52	78.74	69.39	68.64	
Maximum	82.56	82.56	77.47	74.91	

**Table 25: Nursing Care FY 2005/2006** 

Table 25: Natsing Care 11 2000/2000					
	All Hospitals	Small	Community	Teaching	
Valid N	90	18	58	14	
Mean	70.72	79.42	69.86	70.47	
Std Deviation	24.31	15.05	24.72	27.98	
Minimum	58.35	70.92	58.35	64.83	
25 <sup>th</sup> Percentile	69.39	78.25	69.09	68.34	
Median	72.02	81.35	71.10	70.39	
75 <sup>th</sup> Percentile	76.26	83.90	73.59	72.66	
Maximum	85.79	85.79	81.93	76.26	

#### Reporting Results by Sex

The E-Scorecard will include the sex difference values [i.e. (F-M)] for each indicator and an indication of the direction (i.e. F>M or M>F) and the statistical significance of these values at a hospital level. Participating hospitals may access their own and other hospitals' difference values and the direction (i.e. F>M or M>F) and statistical significance of these differences for each indicator on a password-protected database at www.hospitalreport.ca.

The interpretation of these data and notes about suppression will accompany this database. In terms of interpretation, if this value [i.e. (F-M)] is negative, males have higher scores than females; if this value is positive, females have higher scores than males. A value of "0" is used as the benchmark as it represents true equity between women and men. Furthermore, if a hospital's 95% confidence interval around their specific value of the difference between women and men for a given indicator includes zero, then the hospital is said to have no statistically significant sex difference for that indicator (which is preferred). If a hospital's 95% confidence interval around their specific value of the difference between women and men for a given indicator does not include zero and is negative, then the hospital is said to have unequal (i.e. M>F) performance or a statistically significant sex difference, in which males have a higher score than females. If a hospital's 95% confidence interval around their specific value of the difference between women and men for a given indicator does not include zero and is positive, then the hospital is said to have unequal (F>M) performance or a statistically significant sex difference, in which females have a significantly higher score than males.

#### **Summary of Results**

This year, small hospitals generally continue to outperform larger community and teaching hospitals in all four dimensions of patient satisfaction.

Of all the indicators, the lowest average scores across the province were related to responsiveness. However, while this also holds across teaching and community hospitals, the small hospitals scored lowest in the area of communication. In relation to other hospital types, small hospitals appear to be very strong performers in the Responsiveness dimension, where scores are significantly higher, and where nearly all are identified as "above average" performers. Please note that there is considerable variation in performance of all indicators. Therefore, hospitals can use opportunities in different areas to improve patients satisfaction levels.

## **Appendix A: Questionnaire Items**

Questionnaire items, including Hospital Report scoring scheme.

\* Indicates that the item is not part of any of the indicators

#	Question	Response options	Scoring
1	When you arrived at the Emergency Department, did	Yes, completely	100
	the first person who took your information answer	Yes, somewhat	50
	your questions?	No	0
		Didn't Have Questions	no score
2	How would you rate the courtesy of the FIRST	Poor	0
	person who took your information?	Fair	25
		Good	50
		Very Good	75
		Excellent	100
3	After you arrived at the Emergency Department, how	Right Away	100
	long was it until you talked to a NURSE about your	15 minutes or less	50
	illness or injury?	More than 15 minutes	0
		Don't know	no score
4*	Once you went to a bed or an examination room,	Less than ½ hour	
	about how long did you have to wait to see a doctor?	Between ½ hour and 1 hour	
	about non-long and you make to make to be a decient	1 to 2 hours	
		More than 2 hours	
		I did not wait at all	
5	If you had to wait to be seen, did someone from the	Yes	100
5	Emergency Department explain the reason for the	No	0
	delay?	1	_
6	<b>-</b>	Didn't have to wait	no score
6	Did someone in the Emergency Department help get	Yes	100
	your messages to family or friends?	No	0
_		I had no messages	0
7	Was there one particular doctor in charge of your	Yes	100
	care in the Emergency Department?	No	0
		Not Sure	no score
8	Did you have to wait too long to see a doctor?	Yes, definitely	0
		Yes, somewhat	50
		No	100
9	When you had important questions to ask a doctor,	Yes, always	100
	did you get answers you could understand?	Yes, sometimes	50
		No	0
		Didn't have any questions	no score
10	If you had any anxieties or fears about your condition	Yes, completely	100
	or treatment, did a doctor discuss them with you?	Yes, somewhat	50
		No	0
		Didn't have anxieties or fears	no score
11	Did you have confidence and trust in the doctors	Yes, always	100
	treating you?	Yes, sometimes	50
		No	0
12*	Did doctors talk in front of you as if you weren't	Yes, often	
	there?	Yes, sometimes	
		No	
13*	After you had seen a doctor in the Emergency	Yes	
	Department, was another doctor or specialist called	No	
	in to see you?	I did not see a doctor	
14	Did you wait too long for this other doctor or	Yes, definitely	0
• •	specialist?	Yes, somewhat	50
		No	100
		No other doctor was needed	no score
15	How would you rate the courtesy of your doctors?	Poor	0
	House you rate the country of your doctors:	Fair	25
		Good	50
		Very Good	75
10	When you had imported supplier - t!	Excellent	100
16	When you had important questions to ask a nurse,	Yes, always	100
	did you get answers you could understand?	Yes, sometimes	50
		No	0

		Didn't have any questions	no score
17	If you had any anxieties or fears about your condition	Yes, completely	100
	or treatment, did a nurse discuss them with you?	Yes, sometimes	50
		No	0
		Didn't have anxieties or fears	no score
18	Did you have confidence and trust in the nurses	Yes, always	100
	treating you?	Yes, sometimes	50
		No	0
19*	Did nurses talk in front of you as if you weren't	Yes, often	
	there?	Yes, sometimes	
		No	
20	How would you rate the courtesy of your nurses?	Poor	0
		Fair	25
		Good	50
		Very Good	75
		Excellent	100
21	How would you rate the availability of your nurses?	Poor	0
	l lon notice you rate the arandomy or your marcos.	Fair	25
		Good	50
		Very Good	75
		Excellent	100
22*	Did you get any tests (such as blood, urine, or x-	Yes	
	rays) when you visited the Emergency Department?	No (go to #26)	
23	Did you wait too long to get your tests?	Yes, definitely	0
20	Bid you want too long to got your toolo.	Yes, somewhat	50
		No	100
24	Did someone explain why you needed these tests in	Yes, completely	100
27	a way that you could understand?	Yes, somewhat	50
	a way that you could understand:	No	0
25	Did someone explain the results of the tests in a way	Yes, completely	100
23	that you could understand?	Yes, somewhat	50
	that you could understand:	No	0
26*	Were you ever in any pain?	Yes	-
20	Were you ever in any pain:	No (Go to #31)	
27*	When you had pain, was it usually sever, moderate,	Severe	
21	or mild?	Moderate	
	or mild:	Mild	
28*	Did you ever request pain medicine in the	Yes	
20	Emergency Department?	No	
29	Do you think that the Emergency Department staff	Yes, definitely	100
23	did everything they could to help control your pain?	Yes, somewhat	50
	and everything they could to help control your pall?	No	0
30	Overall, how much pain medicine did you get?	Not enough	0
50	overall, now much pain medicine did you get?	Right amount	100
		Too much	0
		I did not get pain medicine	no score
31	Were you told what danger signals about your illness	Yes, completely	100
J1	or injury to watch out for when you got home?	Yes, somewhat	50
	or injury to water out for when you got nome!	No	0
32*	Before you left the Emergency Department, were	Yes	+
JZ	any new medications prescribed or ordered for you?	No	
33	Did someone explain how to take the new	Yes, completely	100
JJ	medications?	Yes, somewhat	50
	medications:	No	0
			-
24	Did compone tell you shout side offeets the	Didn't need explanation	no score
34	Did someone tell you about side effects the	Yes, completely	100
	medicines might have?	Yes, somewhat	50
		No Didn't pood explanation	0
25*	Did you need further treatment of a record of the	Didn't need explanation	no score
35*	Did you need further treatment after you left the	Yes	
	Emergency Department?	No (Go to #37)	

36*	Was an appointment made for this treatment before	Yes, with a new doctor or nurse	
	you left the Emergency Department?	Yes, with the same doctor or	
		nurse	
		No	
37	Did you know who to call if you needed help or had	Yes	100
	more questions after you left the Emergency	No	0
	Department?	Not sure	0
38*	About how long did you spend in the Emergency	Less than 1 hour	
	Department from the time you arrived to the time you	Between 1 and 3 hours	
	left?	Between 3 and 6 hours	
		Between 6 and 10 hours	
		Between 10 and 12 hours	
		Between 12 and 24 hours	
		More than 24 hours	
39	How would you rate the amount of time you spent in	Poor	0
	the Emergency Department?	Fair	25
	and Emergency - Spannism	Good	50
		Very Good	75
		Excellent	100
40	While you were in the Emergency Department, were	Yes, completely	100
-	you able to get all the services you needed?	Yes, somewhat	50
	,	No	0
41	Were the possible causes of your problem explained	Yes, completely	100
''	in a way that you could understand?	Yes, somewhat	50
	in a way that you sould understand.	No	0
		Didn't need explanation	no score
42	While you were in the Emergency Department, were	Yes, often	0
72	there time when you did not get the help you	Yes, sometimes	50
	needed?	No	100
	nodou:	Didn't need help	no score
43	Did each hospital staff person treat you with dignity	Yes, always	100
73	and respect?	Yes, sometimes	50
	and respect.	No	0
44	Did you have enough say about your care?	Yes, definitely	100
	Bid you have chough out about your oute.	Yes, somewhat	50
		No	0
45	Did you feel you had enough privacy during your	Yes, always	100
	Emergency Department visit?	Yes, sometimes	50
	Emergency Department viole:	No	0
		Doesn't apply	no score
46	Overall, how would you rate the care you received in	Poor	0
	the Emergency Department?	Fair	25
		Good	50
		Very Good	75
		Excellent	100
47	How would you rate the courtesy of the Emergency	Poor	0
l	Department staff?	Fair	25
		Good	50
		Very Good	75
		Excellent	100
48	How would you rate the explanation of what was	Poor	0
1	done to you?	Fair	25
		Good	50
		Very Good	75
		Excellent	100
49	How would you rate how well the doctors and nurses	Poor	0
.	worked together?	Fair	25
		Good	50
		Very Good	75
		Excellent	100
50	Would you recommend this Emergency Department	Yes, definitely	100
	jou .cccc.uo Emergency Department	,	

	to family and friends?	Yes, probably No	50 0
51	Was the entire Emergency Department as clean as it should have been?	Yes, definitely Yes, somewhat No	100 50 0
52*	In general, how would you rate your health?	Poor Fair Good Very Good Excellent	
53*	Do you have a regular family physician/general practitioner who you see when you have health problems?	Yes No	
54*	How serious was the injury or illness that prompted you to come to the Emergency Department?	Extremely serious Very serious Moderately serious Slightly serious Not at all serious	
55*	During the past month, how many days did illness or injury keep you in bed all or part of the day?	None One Day Two Days Three Days Four Days Five to Seven Days Eight to Ten Days More than Ten Days	
56*	In the last 6 months, have you been a patient in a hospital overnight or longer?	No Yes, only one time Yes, more than one time	

## Hospital Report 2007: Emergency Department Care

57*	What is the highest grade or level of school that you have completed?	Public school High school College, trade, or technical school University undergraduate degree Post university/graduate education	
58*	Who completed this survey?	Patient Someone else	